


Patent Claims

1. Process for the preparation of compounds of the formula I



(I),

in which


R^1 represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

R^2 represents straight-chain alkyl having up to 4 carbon atoms,

R^3 and R^4 are identical or different and represent a straight-chain or branched alkyl chain having up to 5 carbon atoms, which is optionally substituted up to two times in an identical or different manner by hydroxyl or methoxy,

or

R^3 and R^4 , together with the nitrogen atom, form a piperidinyl, morpholinyl or thiomorpholinyl ring or a radical of the formula



in which

contd.
a¹

5

~~R⁷ denotes hydrogen, formyl, straight-chain or branched acyl or alkoxy carbonyl each having up to 6 carbon atoms, or straight-chain or branched alkyl having up to 6 carbon atoms, which is optionally mono- to disubstituted, in an identical or different manner, by hydroxyl, carboxyl, straight-chain or branched alkoxy or alkoxy carbonyl each having up to 6 carbon atoms, or denotes C₃₋₈-cycloalkyl,~~

10

~~and the heterocycles mentioned under R³ and R⁴, formed together with the nitrogen atom, are optionally mono- to disubstituted, in an identical or different manner, if appropriate also geminally, by hydroxyl, formyl, carboxyl, straight-chain or branched acyl or alkoxy carbonyl each having up to 6 carbon atoms,~~

15

~~and/or the heterocycles mentioned under R³ and R⁴, formed together with the nitrogen atom, are optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms, which is optionally mono- to disubstituted, in an identical or different manner, by hydroxyl or carboxyl,~~

20

~~and/or the heterocycles mentioned under R³ and R⁴, formed together with the nitrogen atom, are optionally substituted by piperidinyl or pyrrolidinyl linked via N,~~

25


~~R⁵ and R⁶ are identical or different and represent hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or straight-chain or branched alkoxy having up to 6 carbon atoms,~~

~~characterized in that compounds of the formula (II)~~

30

10022954-121701


contd.
A¹



in which

R¹, R², R⁵ and R⁶ have the meanings indicated above,

are reacted with sulphuric acid to give compounds of the formula (III)



in which

R¹, R², R⁵ and R⁶ have the meanings indicated above,

and then with thionyl chloride and the product thus obtained is reacted in situ in an inert solvent with an amine of the formula (IV)



in which

R³ and R⁴ have the meaning indicated above,

contd.
a¹

and, if appropriate, reacted to give the corresponding salts, hydrates or N-oxides.

5 2. Process according to Claim 1, characterized in that


R¹ denotes hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

10 R² denotes straight-chain alkyl having up to 4 carbon atoms,

R³ and R⁴ identically to or differently from one another denote a straight-chain or branched alkyl chain having up to 5 carbon atoms, which is optionally substituted up to two times in an identical or different manner by hydroxyl or methoxy,

or

R³ and R⁴, together with the nitrogen atom, form a piperidinyl or morpholinyl ring or a radical of the formula



in which

25 R⁷ denotes hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms, which is optionally mono- or disubstituted, in an identical or different manner, by hydroxyl, straight-chain or branched alkoxy each having up to 4 carbon atoms, or
30 denotes C₃₋₆-cycloalkyl,

10022954-12101

Sub
B²

and the heterocycles mentioned under R^3 and R^4 , formed together with the nitrogen atom, are optionally mono- or disubstituted, in an identical or different manner, if appropriate also geminally, by hydroxyl, straight-chain or branched acyl or alkoxy carbonyl each having up to 4 carbon atoms, optionally by straight-chain or branched alkyl having up to 4 carbon atoms, which is optionally mono- or disubstituted, in an identical or different manner, by hydroxyl,

R^5 and R^6 identically to or differently from one another denote hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or straight-chain or branched alkoxy having up to 6 carbon atoms.

3. Process according to Claim 1, characterized in that


R^1 denotes hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

R^2 denotes straight-chain alkyl having up to 4 carbon atoms,

R^3 and R^4 identically to or differently from one another denote methyl or ethyl, which are optionally substituted up to two times in an identical or different manner by hydroxyl,

or

R^3 and R^4 , together with the nitrogen atom, form a piperidinyl or morpholinyl ring or a radical of the formula



in which

R^7 denotes hydrogen, methyl or ethyl, which is optionally mono- or disubstituted, in an identical or different manner, by hydroxyl, methoxy or ethoxy, or denotes cyclopentyl or cyclohexyl,

and the heterocycles mentioned under R^3 and R^4 , formed together with the nitrogen atom, are optionally mono- or disubstituted, in an identical or different manner, if appropriate also geminally, by hydroxyl, methyl or ethyl,

R^5 and R^6 identically to or differently from one another denote hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or straight-chain or branched alkoxy having up to 6 carbon atoms.

4. Process according to Claim 1, characterized in that

R^1 denotes methyl or ethyl,

R^2 denotes n-propyl,


R^3 and R^4 identically to or differently from one another denote methyl or ethyl, which are optionally substituted up to two times in an identical or different manner by hydroxyl,

or

R^3 and R^4 , together with the nitrogen atom, form a piperidinyll or morpholinyl ring or a radical of the formula

Sub
B2

10022954.121701



in which


R⁷ denotes hydrogen, methyl or ethyl, which is optionally mono- or disubstituted, in an identical or different manner, by hydroxyl, methoxy or ethoxy, or denotes cyclopentyl or cyclohexyl,

and the heterocycles mentioned under R³ and R⁴, formed together with the nitrogen atom, are optionally mono- or disubstituted, in an identical or different manner, if appropriate also geminally, by hydroxyl, methyl or ethyl,

R⁵ denotes hydrogen,

R⁶ denotes ethoxy

5. Process according to Claim 1, characterized in that the compounds of the formula (II) are prepared by reaction of the compounds of the formula (V)



in which

R¹ and R² have the meaning indicated Claim 1


and

Sub
B2

10022954-121701

L represents straight-chain or branched alkyl having up to 4 carbon atoms,

with compounds of the general formula (VI)



in which

R⁵ and R⁶ have the meaning indicated in Claim 1

in a two-stage reaction in the systems methanol and phosphorous oxychloride/
acetic acid or methanol and acetyl chloride / acetic acid.

10022954-121701